Author index

Abbate, E. see *Schierloh, P.*
Addison, E.G., North, J., Bakhsh, I., Marden, C., Haq, S., Al-Sarraj, S., Malayeri, R., Wickremasinghe, R.G., Davies, J.K. and *Lowdell, M.W.* Ligation of CD8α on human natural killer cells prevents activation-induced apoptosis and enhances cytolytic activity, 354
Aho, T.L.T., Lund, R.J., Ylikoski, E.K., Matikainen, S., Lahesmaa, R. and *Koskinen, P.J.* Expression of human *pim* family genes is selectively up-regulated by cytokines promoting T helper type 1, but not T helper type 2, cell differentiation, 82
Akamatsu, T. see *Yao, C.*
Akita, Y. see *Uchida, K.*
Akiyama, Y. see *Kaneko, M.*
Alemán, M. see *Schierloh, P.*
de Alencar, R. see *Nomizo, A.*
Aliesky, H. see *Kala, M.*
Al-Sarraj, S. see *Addison, E.G.*
Alves, L. see *Schierloh, P.*
Ando, M. see *Nishijima, K.-i.*
Arikuni, H. see *Takata-Tomokuni, A.*

Bakhsh, I. see *Addison, E.G.*
Balk, S.P. see *Sonoda, K.-H.*
Bamford, K.B. see *Scott, K.*
de la Barrera, S. see *Schierloh, P.*
Barth, P.J. see *Steiniger, B.*
Bates, J.T. and *Bucy, R.P.* Enhanced responsiveness to antigen contributes more to immunological memory in CD4 T cells than increases in the number of cells, 318
Battaglia, A. see *Fattorossi, A.*
Bellinghausen, I., König, B., Böttcher, I., Knop, J. and *Saloga, J.* Regulatory activity of human $CD4^+$ $CD25^+$ T cells depends on allergen concentration, type of allergen and atopy status of the donor, 103
Biedroń, R. see *Jozefowski, S.*
Bobek, M. see *Jozefowski, S.*
de Bono, B. see *Rhodes, D.A.*
Borie, D.C. see *Kleihauer, A.*
Borron, P.J. see *Seabrook, T.J.*
Bose, B. and *Sinha, S.* Problems in using statistical analysis of replacement and silent mutations in antibody genes for determining antigen-driven affinity selection, 172
Böttcher, I. see *Bellinghausen, I.*
Brito, G.A.C. see *Nascimento, S.B.*
Brown, C.B. see *Unruh, T.L.*
Bucy, R.P. see *Bates, J.T.*
Buzzonetti, A. see *Fattorossi, A.*

Canada, N. see *Teixeira, L.*
Cardillo, F. see *Nomizo, A.*
Carosella, E.D. see *Rond, S.L.*
Chapman, H.A. see *Kala, M.*
Chen, C.-R. see *Kala, M.*
Chiang, A.K.S. see *Wong, O.H.*
Cho, D. see *Park, J.*
Ciaraffa, F. see *Fattorossi, A.*
Cooke, A. see *Phillips, J.M.*
da Costa, J.M.C. see *Teixeira, L.*
Cunha, F.Q. see *Nascimento, S.B.*

Dai, J. see *Li, B.*
Dalpke, A.H., Lehner, M.D., Hartung, T. and *Heeg, K.* Differential effects of CpG-DNA in Toll-like receptor-2/-4/-9 tolerance and cross-tolerance, 203
Davies, J.K. see *Addison, E.G.*
Deans, J.P. see *Unruh, T.L.*
Dessi, D. see *Scott, K.*
Dieleman, L.A. see *Qian, B.-F.*
Ding, A., Yu, H., Yang, J., Shi, S. and *Ehrt, S.* Induction of macrophage-derived SLPI by *Mycobacterium tuberculosis* depends on TLR2 but not MyD88, 381
Downing, J.E.G. see *Shepherd, A.J.*
Dudler, L. see *Seabrook, T.J.*

Ehrt, S. see *Ding, A.*
Ellis, R.J., Varela-Calvino, R., Tree, T.I.M. and *Peakman, M.* HLA Class II molecules on haplotypes associated with type 1 diabetes exhibit similar patterns of binding affinities for coxsackievirus P2C peptides, 337
Erlandsen, S.L. see *Sarkar, K.*
Evoli, A. see *Fattorossi, A.*
Exley, M. see *Sonoda, K.-H.*

Fan, Y. see *Qiu, H.*
Fattorossi, A., Battaglia, A., Buzzonetti, A., Ciaraffa, F., Scambia, G. and *Evoli, A.* Circulating and thymic $CD4^+$ $CD25^+$ T regulatory cells in myasthenia gravis: effect of immunosuppressive treatment, 134
Faustino, A.M.R. see *Teixeira, L.*
Felföldi, B., Imre, G., Igyártó, B., Iván, J., Mihalik, R., Lackó, E., Oláh, I. and *Magyar, A. In ovo* vitelline duct ligation results in transient changes of bursal microenvironments, 267
Ferreira, P. see *Teixeira, L.*
Fiori, P.L. see *Scott, K.*
Foster, G.R. see *Scott, K.*
Fujimoto, W. see *Takata-Tomokuni, A.*
Fujiwara, S. see *Uchida, K.*

Germain, C. see *Scott, K.*
Gockel, C.M. and *Russell, M.W.* Induction and recall of immune memory by mucosal immunization with a non-toxic recombinant enterotoxin-based chimeric protein, 477

Gomes, A.S. see *Nascimento, S.B.*
Gonzalez, A. see *Rond, S.L.*
Gonzalez, A.S.L. see *Rond, S.L.*
González-Amaro, R., Mittelbrunn, M. and *Sánchez-Madrid, F.* Therapeutic anti-integrin (α4 and αL) monoclonal antibodies: two-edged swords?, 289
Gorgone, D.A. see *Jackson, S.S.*
Gregory, C.R. see *Kleihauer, A.*
Grigoriou, L. see *Unruh, T.L.*
Guérin, M. see *Néron, S.*
Guerrant, R.L. see *Nascimento, S.B.*
Guo, Y. see *Li, B.*

Hachiya, H. see *Uchida, K.*
Han, X. see *Qiu, H.*
Hanlon, A., Jang, S. and *Salgame, P.* Cbl-b differentially regulates activation-induced apoptosis in T helper 1 and T helper 2 cells, 507
Haq, S. see *Addison, E.G.*
Hartung, T. see *Dalpke, A.H.*
Haskins, K. see *Phillips, J.M.*
Hatayama, T. see *Takata-Tomokuni, A.*
Hay, J.B. see *Seabrook, T.J.*
Hayashi, Y. see *Kato, M.*
Hayashi-Ozawa, A. see *Nishijima, K.-i.*
Hedrich, H.-J. see *Milićević, N.M.*
Heeg, K. see *Dalpke, A.H.*
Heike, Y. see *Ikarashi, Y.*
Hisatomi, T. see *Sonoda, K.-H.*
Hoentjen, F. see *Qian, B.-F.*
Hopkins, J. see *Mohan, J.*
Hosoi, K. see *Yao, C.*
Hou, S. see *Li, B.*
Huang, F.-P. see *Wong, O.H.*
Hultkrantz, S., Östman, S. and *Telemo, E.* Induction of antigen-specific regulatory T cells in the liver-draining celiac lymph node following oral antigen administration, 362
Hyodoh, F. see *Takata-Tomokuni, A.*

Igyártó, B. see *Felföldi, B.*
Iijima, S. see *Nishijima, K.-i.*
Iizuka, A. see *Ikarashi, Y.*
Ikarashi, Y., Iizuka, A., Koshidaka, Y., Heike, Y., Takaue, Y., Yoshida, M., Kronenberg, M. and *Wakasugi, H.* Phenotypical and functional alterations during the expansion phase of invariant Vα14 natural killer T (Vα14i NKT) cells in mice primed with α-galactosylceramide, 30
Imre, G. see *Felföldi, B.*
Ishibashi, T. see *Sonoda, K.-H.*
Isozaki, Y. see *Takata-Tomokuni, A.*
Iván, J. see *Felföldi, B.*
Izumi, T. see *Kato, M.*

Jackson, S.S., Schmitz, J.E., Kuroda, M.J., McKay, P.F., Sumida, S.M., Martin, K.L., Yu, F., Lifton, M.A., Gorgone, D.A. and *Letvin, N.L.* Evaluation of CD62L expression as a marker for vaccine-elicited memory cytotoxic T lymphocytes, 443
Jacob, R. see *Steiniger, B.*
Jang, S. see *Hanlon, A.*
Jeganathan, A. see *Reed, A.J.*
Ji, J. see *Li, B.*
Jiao, L. see *Qiu, H.*
Jones, M. see *Scott, K.*
Joyee, A.G. see *Qiu, H.*
Jozefowski, S., Biedroń, R., Bobek, M. and *Marcinkiewicz, J.* Leukotrienes modulate cytokine release from dendritic cells, 418

Kala, M., Chen, C.-R., McLachlan, S.M., Rapoport, B., Aliesky, H. and *Chapman, H.A.* Cathepsin S is not crucial to TSHR processing and presentation in a murine model of Graves' disease, 532
Kanamori, N. see *Yao, C.*
Kaneko, M., Akiyama, Y., Takimoto, H. and *Kumazawa, Y.* Mechanism of up-regulation of immunoglobulin A production in the intestine of mice unresponsive to lipopolysaccharide, 64
Katagiri, K. see *Kinashi, T.*
Kato, M., Yamaguchi, T., Tachibana, A., Suzuki, M., Izumi, T., Maruyama, K., Hayashi, Y. and *Kimura, H.* An atypical protein kinase C, PKC ζ, regulates human eosinophil effector functions, 193
Katsuyama, H. see *Takata-Tomokuni, A.*
Kawai, T. see *Uchida, K.*
Kazaoka, Y. see *Uchida, K.*
Kearns-Jonker, M. see *Kleihauer, A.*
Kim, M.S. and *Sin, J.-I.* Both antigen optimization and lysosomal targeting are required for enhanced anti-tumour protective immunity in a human papillomavirus E7-expressing animal tumour model, 255
Kim, S.H. see *Park, J.*
Kim, T.S. see *Park, J.*
Kimura, H. see *Kato, M.*
Kinashi, T. and *Katagiri, K.* Regulation of immune cell adhesion and migration by regulator of adhesion and cell polarization enriched in lymphoid tissues, 164
Kinoshita, Y. see *Nishijima, K.-i.*
Kleihauer, A., Gregory, C.R., Borie, D.C., Kyles, A.E., Shulkin, I., Patanwala, I., Zahorsky-Reeves, J., Starnes, V.A., Mullen, Y., Todorov, I.T. and *Kearns-Jonker, M.* Identification of the V_H genes encoding xenoantibodies in non-immunosuppressed rhesus monkeys, 89
Knop, J. see *Bellinghausen, I.*
König, B. see *Bellinghausen, I.*
Koshidaka, Y. see *Ikarashi, Y.*
Koskinen, P.J. see *Aho, T.L.T.*
Kosugi-Tanaka, C. see *Yao, C.*
Kozaki, K.-i. see *Uchida, K.*
Kronenberg, M. see *Ikarashi, Y.*
Kruhlak, M.J. see *Sarkar, K.*
Kumazawa, Y. see *Kaneko, M.*
Kuroda, J. see *Muto, S.-i.*
Kuroda, M.J. see *Jackson, S.S.*
Kusaka, M. see *Takata-Tomokuni, A.*
Kyles, A.E. see *Kleihauer, A.*

Lackó, E. see *Felföldi, B.*
Lahesmaa, R. see *Aho, T.L.T.*
Lechler, R.I. see *Scott, K.*
Lehner, M.D. see *Dalpke, A.H.*
Letvin, N.L. see *Jackson, S.S.*
Li, B., Wang, H., Dai, J., Ji, J., Qian, W., Zhang, D., Hou, S. and *Guo, Y.* Construction and characterization of a humanized anti-human CD3 monoclonal antibody 12F6 with effective immunoregulation functions, 487
Li, H. see *Unruh, T.L.*
Li, X. see *Yao, C.*
Lifton, M.A. see *Jackson, S.S.*
Lombardi, G. see *Scott, K.*
Lowdell, M.W. see *Addison, E.G.*
Lund, R.J. see *Aho, T.L.T.*

Mabbott, N.A. see *Mohan, J.*
Madureira, P. see *Teixeira, L.*
Magyar, A. see *Felföldi, B.*
Malayeri, R. see *Addison, E.G.*
Manunta, M. see *Scott, K.*
Marcinkiewicz, J. see *Jozefowski, S.*
Marden, C. see *Addison, E.G.*
Marques, A. see *Teixeira, L.*
Martin, K.L. see *Jackson, S.S.*
Martins, M.J.B. see *Nascimento, S.B.*
Maruyama, K. see *Kato, M.*
Matikainen, S. see *Aho, T.L.T.*
Matsumoto, K. see *Muto, S.-i.*
Matsumoto, Y. see *Uchida, K.*
Matsuo, K. see *Uchida, K.*
McKay, P.F. see *Jackson, S.S.*
McLachlan, S.M. see *Kala, M.*
Meireles, C.S. see *Teixeira, L.*
Mengel, J. see *Nomizo, A.*
Mihalik, R. see *Felföldi, B.*
Milićević, N.M., Nohroudi, K., Milićević, Ž., Hedrich, H.-J. and *Westermann, J.* T cells are required for the peripheral phase of B-cell maturation, 308
Milićević, Ž. see *Milićević, N.M.*
Mitchell, P. see *Scott, K.*
Mittelbrunn, M. see *González-Amaro, R.*
Miyan, J.A. see *Shepherd, A.J.*
Mohan, J., Hopkins, J. and *Mabbott, N.A.* Skin-derived dendritic cells acquire and degrade the scrapie agent following *in vitro* exposure, 122
Monnai, M. see *Muto, S.-i.*
Mullen, Y. see *Kleihauer, A.*
Murakami, M. see *Muto, S.-i.*
Murdiastuti, K. see *Yao, C.*
Mutch, C.M. see *Unruh, T.L.*
Muto, S.-i., Monnai, M., Okuhara, Y., Murakami, M., Kuroda, J., Ono, T. and *Matsumoto, K.* Altered cytokine expression in mesenteric lymph nodes in a rat strain (Matsumoto Eosinophilic Shinshu) that spontaneously develops hypereosinophilia, 373

Naganawa, Y. see *Uchida, K.*
Naji, A. see *Reed, A.J.*
Nakagawa, A. see *Uchida, K.*
Nascimento, S.B., Sousa, R.B., Martins, M.J.B., Gomes, A.S., Souza, M.H.L.P., Guerrant, R.L., Cunha, F.Q., Ribeiro, R.A. and *Brito, G.A.C.* Glutamine depletion potentiates leucocyte-dependent inflammatory events induced by carrageenan or *Clostridium difficile* toxin A in rats, 328
Néron, S., Racine, C., Roy, A. and *Guérin, M.* Differential responses of human B-lymphocyte subpopulations to graded levels of CD40–CD154 interaction, 454
Nishijima, K.-i., Ando, M., Sano, S., Hayashi-Ozawa, A., Kinoshita, Y. and *Iijima, S.* Costimulation of T-cell proliferation by anti-L-selectin antibody is associated with the reduction of a cdk inhibitor p27, 347
Nohroudi, K. see *Milićević, N.M.*
Nombela-Arrieta, C. see *Stein, J.V.*
Nomizo, A., Postol, E., de Alencar, R., Cardillo, F. and *Mengel, J.* Natural killer T cells are required for the development of a superantigen-driven T helper type 2 immune response in mice, 233
Noorchashm, H. see *Reed, A.J.*
North, J. see *Addison, E.G.*

Oh-iwa, I. see *Uchida, K.*
Ohura, K. see *Uchida, K.*
Okuhara, Y. see *Muto, S.-i.*
Oláh, I. see *Felföldi, B.*
Ono, T. see *Muto, S.-i.*
Oshima, T. see *Sonoda, K.-H.*
Ostergaard, H.L. see *Shen, A.*
Östman, S., Taube, M. and *Telemo, E.* Tolerosome-induced oral tolerance is MHC dependent, 464
Östman, S. see *Hultkrantz, S.*
Otsuki, T. see *Takata-Tomokuni, A.*

Park, J., Kim, S.H., Cho, D. and *Kim, T.S.* Formononetin, a phyto-oestrogen, and its metabolites up-regulate interleukin-4 production in activated T cells via increased AP-1 DNA binding activity, 71
Patanwala, I. see *Kleihauer, A.*
Peakman, M. see *Ellis, R.J.*
Perate, A.L. see *Reed, A.J.*
Phillips, J.M., Haskins, K. and *Cooke, A.* MAdCAM-1 is needed for diabetes development mediated by the T cell clone, BDC-2·5, 525
Postol, E. see *Nomizo, A.*
Puente, L.G. see *Shen, A.*

Qian, B.-F., Tonkonogy, S.L., Hoentjen, F., Dieleman, L.A. and *Sartor, R.B.* Dysregulated luminal bacterial antigen-specific T-cell responses and antigen-presenting cell function in HLA-B27 transgenic rats with chronic colitis, 112
Qian, W. see *Li, B.*
Qiao, H. see *Sonoda, K.-H.*
Qiu, H., Wang, S., Yang, J., Fan, Y., Joyee, A.G., Han, X., Jiao, L. and *Yang, X.* Resistance to chlamydial lung infection is dependent on major histocompatibility complex as well as non-major histocompatibility complex determinants, 499

Racine, C. see *Néron, S.*
Rapoport, B. see *Kala, M.*

Reed, A.J., Zarrabi, Y., Perate, A.L., Jeganathan, A., Naji, A. and *Noorchashm, H.* The frequency of double-positive thymocytes expressing an αβ TCR clonotype regulates peripheral CD4 T cell compartment homeostasis, 400
Rhodes, D.A., de Bono, B. and *Trowsdale, J.* Relationship between SPRY and B30.2 protein domains. Evolution of a component of immune defence?, 411
Ribeiro, A. see *Teixeira, L.*
Ribeiro, R.A. see *Nascimento, S.B.*
Rodrigues, D. see *Teixeira, L.*
Roldán, N. see *Schierloh, P.*
Rond, S.L., Gonzalez, A., Gonzalez, A.S.L., Carosella, E.D. and *Rouas-Freiss, N.* Indoleamine 2,3 dioxygenase and human leucocyte antigen-G inhibit the T-cell alloproliferative response through two independent pathways, 297
Rouas-Freiss, N. see *Rond, S.L.*
Roy, A. see *Néron, S.*
Russell, M.W. see *Gockel, C.M.*

Saga, S. see *Uchida, K.*
Sakamoto, T. see *Sonoda, K.-H.*
Salgame, P. see *Hanlon, A.*
Saloga, J. see *Bellinghausen, I.*
Sánchez-Madrid, F. see *González-Amaro, R.*
Sano, S. see *Nishijima, K.-i.*
Sanyal, R. see *Unruh, T.L.*
Sarkar, K., Kruhlak, M.J., Erlandsen, S.L. and *Shaw, S.* Selective inhibition by rottlerin of macropinocytosis in monocyte-derived dendritic cells, 513
Sartor, R.B. see *Qian, B.-F.*
del C. Sasiain, M. see *Schierloh, P.*
Scambia, G. see *Fattorossi, A.*
Schierloh, P., Alemán, M., Yokobori, N., Alves, L., Roldán, N., Abbate, E., del C. Sasiain, M. and *de la Barrera, S.* NK cell activity in tuberculosis is associated with impaired CD11a and ICAM-1 expression: a regulatory role of monocytes in NK activation, 541
Schmitz, J.E. see *Jackson, S.S.*
Scott, K., Manunta, M., Germain, C., Smith, P., Jones, M., Mitchell, P., Dessi, D., Bamford, K.B., Lechler, R.I., Fiori, P.L., Foster, G.R. and *Lombardi, G.* Qualitatively distinct patterns of cytokines are released by human dendritic cells in response to different pathogens, 245
Seabra, A.R. see *Teixeira, L.*
Seabrook, T.J., Borron, P.J., Dudler, L., Hay, J.B. and *Young, A.J.* A novel mechanism of immune regulation: interferon-γ regulates retention of $CD4^+$ T cells during delayed type hypersensitivity, 184
Shariat, N. see *Unruh, T.L.*
Shaw, S. see *Sarkar, K.*
Shen, A., Puente, L.G. and *Ostergaard, H.L.* Tyrosine kinase activity and remodelling of the actin cytoskeleton are co-temporally required for degranulation by cytotoxic T lymphocytes, 276
Shepherd, A.J., Downing, J.E.G. and *Miyan, J.A.* Without nerves, immunology remains incomplete – *in vivo veritas*, 145
Shi, S. see *Ding, A.*
Shimozato, K. see *Uchida, K.*
Shiwa, M. see *Takata-Tomokuni, A.*
Shulkin, I. see *Kleihauer, A.*
Silva, C. see *Teixeira, L.*
Sin, J.-I. see *Kim, M.S.*
Sinha, S. see *Bose, B.*
Smith, P. see *Scott, K.*
Sonoda, K.-H., Sakamoto, T., Qiao, H., Hisatomi, T., Oshima, T., Tsutsumi-Miyahara, C., Exley, M., Balk, S.P., Taniguchi, M. and *Ishibashi, T.* The analysis of systemic tolerance elicited by antigen inoculation into the vitreous cavity: vitreous cavity-associated immune deviation, 390
Sousa, R.B. see *Nascimento, S.B.*
Souza, M.H.L.P. see *Nascimento, S.B.*
Starnes, V.A. see *Kleihauer, A.*
Stein, J.V. and *Nombela-Arrieta, C.* Chemokine control of lymphocyte trafficking: a general overview, 1
Steiniger, B., Timphus, E.-M., Jacob, R. and *Barth, P.J.* $CD27^+$ B cells in human lymphatic organs: re-evaluating the splenic marginal zone, 429
Sumida, S.M. see *Jackson, S.S.*
Suzuki, M. see *Kato, M.*

Tachibana, A. see *Kato, M.*
Takata-Tomokuni, A., Ueki, A., Shiwa, M., Isozaki, Y., Hatayama, T., Katsuyama, H., Hyodoh, F., Fujimoto, W., Ueki, H., Kusaka, M., Arikuni, H. and *Otsuki, T.* Detection, epitope-mapping and function of anti-Fas autoantibody in patients with silicosis, 21
Takaue, Y. see *Ikarashi, Y.*
Takimoto, H. see *Kaneko, M.*
Taniguchi, M. see *Sonoda, K.-H.*
Taube, M. see *Östman, S.*
Teixeira, L., Marques, A., Meireles, C.S., Seabra, A.R., Rodrigues, D., Madureira, P., Faustino, A.M.R., Silva, C., Ribeiro, A., Ferreira, P., da Costa, J.M.C., Canada, N. and *Vilanova, M.* Characterization of the B-cell immune response elicited in BALB/c mice challenged with *Neospora caninum* tachyzoites, 38
Telemo, E. see *Östman, S.*
Telemo, E. see *Hultkrantz, S.*
Timphus, E.-M. see *Steiniger, B.*
Todorov, I.T. see *Kleihauer, A.*
Tonkonogy, S.L. see *Qian, B.-F.*
Tree, T.I.M. see *Ellis, R.J.*
Trowsdale, J. see *Rhodes, D.A.*
Tsutsumi-Miyahara, C. see *Sonoda, K.-H.*

Uchida, K., Akita, Y., Matsuo, K., Fujiwara, S., Nakagawa, A., Kazaoka, Y., Hachiya, H., Naganawa, Y., Oh-iwa, I., Ohura, K., Saga, S., Kawai, T., Matsumoto, Y., Shimozato, K. and *Kozaki, K.-i.* Identification of specific autoantigens in Sjögren's syndrome by SEREX, 53
Ueki, A. see *Takata-Tomokuni, A.*
Ueki, H. see *Takata-Tomokuni, A.*
Unruh, T.L., Li, H., Mutch, C.M., Shariat, N., Grigoriou, L., Sanyal, R., Brown, C.B. and *Deans, J.P.* Cholesterol depletion inhibits src family kinase-dependent calcium mobilization and apoptosis induced by rituximab crosslinking, 223

Varela-Calvino, R. see *Ellis, R.J.*
Vilanova, M. see *Teixeira, L.*

Wakasugi, H. see *Ikarashi, Y.*
Wang, H. see *Li, B.*
Wang, S. see *Qiu, H.*
Westermann, J. see *Milićević, N.M.*
Wickremasinghe, R.G. see *Addison, E.G.*
Wong, O.H., Huang, F.-P. and *Chiang, A.K.S.* Differential responses of cord and adult blood-derived dendritic cells to dying cells, 13

Yamaguchi, T. see *Kato, M.*
Yang, J. see *Ding, A.*
Yang, J. see *Qiu, H.*
Yang, X. see *Qiu, H.*
Yao, C., Li, X., Murdiastuti, K., Kosugi-Tanaka, C., Akamatsu, T., Kanamori, N. and *Hosoi, K.* Lipopolysaccharide-induced elevation and secretion of interleukin-1β in the submandibular gland of male mice, 213
Ylikoski, E.K. see *Aho, T.L.T.*
Yokobori, N. see *Schierloh, P.*
Yoshida, M. see *Ikarashi, Y.*
Young, A.J. see *Seabrook, T.J.*
Yu, F. see *Jackson, S.S.*
Yu, H. see *Ding, A.*

Zahorsky-Reeves, J. see *Kleihauer, A.*
Zarrabi, Y. see *Reed, A.J.*
Zhang, D. see *Li, B.*

Subject index

adhesion, 193
allergy, 71, 103
alpha gal carbohydrate, 89
antibodies, 38
antibody responses, 289
anti-Fas autoantibody, 21
antigen
  presentation, 337, 513, 532
  specific cell frequency, 318
antigen-presenting cells, 112
AP-1, 71
apoptosis, 13, 223, 354, 507
autoantibodies, 373
autoimmune
  disease, 134
  diseases, 21
autoimmunity, 337

B cells, 38, 223, 267
B30.2 domain, 411
B-cell
  maturation, 308
  surface molecules, 308
B-cell/T-cell interactions, 308
BDC-2·5, 525
bursa, 267

calcium, 223
Cbl-b, 507
CD25, 103
CD138, 454
CD27, 454
CD28, 507
CD3, 487
CD8, 525
CD40, 245
cdk inhibitor, 347
CD40–CD154, 454
$CD4^+$, 318
$CD27^+$ B cells, 429
$CD8^+$ T lymphocyte, 443
$CD4^+$ T lymphocytes, 112
celiac lymph node, 362
cell surface molecule, 443
cellular
  activation, 30, 276
  differentiation, 400
cervical cancer, 255
chemokines, 1
chicken, 267
Chlamydia trachomatis, 499
cholera toxin, 64, 477
codon optimization, 255
complement 5a, 193
congenic mice, 499
contact sensitivity, 145
cord blood, 13
corticosteroids, 134
costimulation, 347, 507
CpG-DNA, 203
cysteine proteases, 532
cytokine, 373
cytokines, 13, 82
cytotoxicity, 354, 541

degranulation, 193
delayed hypersensitivity, 184
dendritic
  cell, 13
  cells, 103, 122, 164, 245, 267, 418, 513
diabetes, 337, 525
DNA vaccine, 255

enteric bacterial antigens, 112
eosinophilia, 373
eosinophils, 193
evolution, 411
exosome, 464
eye, 390

first dose syndrome, 487
fluid-phase endocytosis, 513

genetics, 289
glutamine, 328
glycolipid, 30
GM-CSF, 513
Graves' hyperthyroidism, 532

haemopoiesis, 145
HIV, 411
HLA-B27 transgenic rats, 112
HLA-G, 297
host defence, 499
human
  B lymphocytes, 454
  lymph node, 429
  papillomavirus, 255
  regulatory T cells (Treg), 103
  spleen, 429
  tonsil, 429
Humanization, 487

IDO, 297
IFI16, 53
IgA, 64
IgV_H gene, 89
IL-4, 513
IL-6, 418
IL-10, 418
IL-12, 418
immunoglobulin
  G, 233
  variable regions, 172
immunoglobulins, 289
immunological memory, 318
immunosuppression, 487
innate immunity, 411
integrin activation, 1
interferon-γ, 30, 464
interleukin-5, 373
interleukin-4, 30, 71
interleukin-10, 245
interleukin-12, 245
interleukin-1β, 213, 328
intestinal epithelial cell, 464

KLHL12, 53
KLHL7, 53

Langerhans cells, 122
leukaemia/lymphoma/myeloma, 223
leukotriene, 418
LFA-1, 164
lipopolysaccharide, 203, 213
lipoteichoic acid, 203
L-selectin, 347
lymph node, 184
lymphatics, 184
lymphoblastoid cells, 13
lymphocyte recirculation, 184
lymphocytes, 1, 164
lysosomal targeting, 255

macrophage, 381
MAdCAM-1, 525
major histocompatibility complex, 337
marginal zone, 429
memory, 145, 443
  cells, 477
MHC, 499
  class II, 464

© 2005 Blackwell Publishing Ltd,

migration, 1
mixed lymphocyte reactions, 297
monoclonal antibody, 487
monocytes, 513, 541
mouse, 443
mucosal, 64
  immunity, 477
mycobacteria, 381

natural killer
  cells, 82, 354, 541
  T cells, 233, 390
necrosis, 13
neonates, 525
Neospora caninum, 38
nerve fibres, 145
neuroimmunology, 145
neutrophil migration, 328

oral tolerance, 362, 464
ovalbumin, 362

p27, 347
pathogens, 245
Peyer's patch, 64
phyto-oestrogen, 71
Pim kinases, 82
platelet-activating factor, 193
protease inhibitor, 381
protein
  kinase C, 193
  kinases/phosphatases, 276

rafts, 507
Rap1, 164
RAPL, 164
rat, 373
receptors, 541
regulatory T cells, 362
repertoire development, 400
rodent, 381

scrapie, 122
secondary lymphoid organs, 1
sequence analysis, 172
SEREX, 53
signal transduction, 276, 381
silicosis, 21
siRNA, 21
Sjögren's syndrome, 53
skin, 122
somatic mutation, 172
staphylococcal enterotoxin B, 233
Streptococcus, 477
submandibular gland, 213
superantigens, 233
superoxide anion, 193

T
  cell, 71
  helper cells, 82
  lymphocytes, 30, 400, 513
  regulatory cells, 134
T-cell proliferation, 347
Th1/Th2, 103
  cells, 507
thymus, 134, 400
tolerance, 203, 233, 390
Toll-like
  receptor-4, 64, 213
  receptors, 203
transforming growth factor-β, 82
transmissible spongiform encephalopathy, 122
tuberculosis, 541
tumour
  immunity, 255
  immunotherapy, 354
  necrosis factor-α, 328

vaccination, 443
viruses, 337
vitelline duct, 267
vitreous cavity, 390

xenoantibody response, 89
xenotransplantation, 289